S. Saffarzadeh¹, A. Jamshidi^{1,*}, A.Hadi-Vencheh¹

¹Department of Mathematics, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran

Abstract

So far, many ways have been provided to solve Multiple Criteria Decision Making (MCDM) problems with interval numbers. Most of these methods rank the alternatives according to two criteria, that is, being close to the positive ideal solution (PIS) and far away from the negative ideal solution (NIS). In this paper, a method is presented for solving MCDM problems with interval numbers, such that being close to PIS and being away from NIS have the same effect in alternatives ranking. In the proposed method, the first PIS and NIS are determined as interval numbers and distance of each alternative from PIS and NIS is calculated by extension of Euclidean distance. Then, a compromise index is defined to rank the alternatives. Three numerical examples are given to compare the proposed method with other methods presented in the literature.

Keywords: Multiple Criteria Decision Making; interval number; criterion weight; positive ideal solution; negative ideal solution.

1. Introduction

Multiple Criteria Decision Making (MCDM) problems are an important research field in decision science, management, and operations research. In MCDM problems, a group of alternatives is evaluated and compared based on their performance under several criteria. Based on this assessment, alternatives are ranked and the alternative which has the most aggregated performance is selected for implementation. There are many methods to solve MCDM problems [1-6]. The classical methods to solve MCDM problems are: Simple Additive Weighting (SAW) method [7] which uses a simple aggregate function, the Analytical Hierarchy Process (AHP) [1] which is based on the judgments of Decision Makers (DMs) to decompose a complex problem into a hierarchy with the goal at the top level of hierarchy; Technique for Order Preference by Similarity to Ideal Solution (TOPSIS) [2] and Vlse Kriterijumska Optimizacija I Kompromisno Resenje (VIKOR) [8] seeking for an alternative -a compromise solution- that is close to Positive Ideal Solution (PIS) as possible and as far away from Negative Ideal Solution (NIS) as possible; Complex Proportional Assessment (COPRAS) [3] technique is initiated by Zavadskas et al. and works similar to SAW; ELimination Et Choix Traduisant la REalité (ELECTRE) method has different types and the study of Govindan and Jespen [5] can be mentioned as a comprehensive source of ELECTRE and ELECTRE-based

* Correspondence: Phone: +98 31 35354001; Mobile: +98 9177049067; Fax: +98 31 35354060

E-mail addresses: alijamshidi63@yahoo.com, s.saffarzadeh@gmail.com, ahadi@khuif.ac.ir

methods. Carneiro et al. [6] consider the effect of cognitive aspects in MCDM. For this purpose, they take credibility, expertise level and behavior style of DMs into account and propose a method that links these aspects to the problem. Abdollahi et al. [9] proposed a method based on the TOPSIS and entropy to rank multifarious Demand Response Resources.

The above-mentioned classical methods have considered the cases that decision matrix includes exact data. But it is not always the case, indeed in most real world problems decision information is expressed as fuzzy or interval numbers. To tackle uncertainty in MCDM problems, an integrated MCDM model based on Fuzzy AHP (FAHP), TOPSIS and grey relational analysis (GRA) in paper industry proposed by Kirubakaran and Ilangkumaran [10]. The authors employ FAHP to determine criteria weights and the alternatives are then ranked by TOPSIS-GRA. Singh et al. [11] proposed an integrated model based on FAHP and FTOPSIS for a selection of third party logistics (3PL) selection in cold chain management. In their method, authors applied FAHP to rank criteria and then they used FTOPSIS to select the best 3PL. Mahmoudi et al. [12] have presented an extension of fuzzy VIKOR to solve Supplier Selection Problem. They proposed a fuzzy distance measure to rank the suppliers. In their method the alternatives are ranked using the preference ratio method. Hu et al. [13] proposed an MCDM problem with stochastic and intuitionistic data.

Although in uncertain MCDM problems, focusing on interval numbers has been less than fuzzy numbers, but useful researches have been done in this case. Jahanshahloo et al. [14] developed the TOPSIS method for MCDM problems with interval numbers. In their research firstly the PIS and NIS are determined by a special algorithm, then the separation of each alternative from the PIS and NIS is calculated using interval arithmetic and finally, alternatives are ranked by using the relative closeness of each alternative with respect to the PIS. Jahanshahloo et al. [15] also extended the TOPSIS method for MCDM problems with interval numbers. They first identify the PIS and NIS of each alternative, then the separation of each alternative from its PIS and NIS is calculated and alternatives are ranked by using relative closeness of each alternative with respect to its PIS. Sayadi et al. [16] developed the VIKOR method for MCDM problems with interval numbers. In their method, first the PIS and NIS are determined, then utility and regret measures and VIKOR index are obtained for each alternative by using interval arithmetic and finally, VIKOR indexes which are interval numbers were compared together. For this purpose, a coefficient called optimism level of DM was introduced and by using it, interval numbers were compared and alternatives were ranked. Dymova et al. [17] developed the TOPSIS method for MCDM problems with interval numbers. Their proposed method is fundamentally different from similar methods. The first difference is that the interval numbers are compared then the PIS and NIS are determined. The second difference is that the separation of each alternative from the PIS and NIS is obtained by calculating the separation between centers of intervals. Using this measure does not require Euclidean distance or Hamming distance to calculate the separation of each alternative from the PIS and NIS and the separation of each alternative from the PIS and NIS is easier to calculate. Hafezolkotob et al. [18] developed the multi MOORA method for MCDM problems with

interval numbers. In their method, first the decision matrices are normalized and weighted, then in order to rank the alternatives and find the best one, the interval numbers are compared by using their degree of possibility. Jahan and Edwards [19] presented a VIKOR method for materials ranking with the simultaneous availability of interval data and all types of criteria.

In this paper, to solve interval MCDM problems a new method is presented. In the proposed method, being close to PIS and being away from NIS have the same effect in alternatives ranking. In the other words, we look for a compromise solution, which is close to the PIS as possible and remotest from the NIS as possible simultaneously. For this purpose, first the initial decision matrix is normalized and the PIS and the NIS determined in interval form. Then, the distance of each alternative from the PIS and the NIS is calculated by extension of Euclidean distance and an index called relative agreement is obtained for each alternative. In the end, the alternatives are ranked by sorting these ratios in descending order.

The rest of this paper is organized as follows. In section 2 some interval definitions and interval arithmetic are given. Section 3 is devoted to the proposed method and its algorithm. In section 4, the proposed method is compared with the interval TOPSIS method. In section 5, three numerical examples are presented to compare the proposed method and those published in the literature. Section 6 is devoted to the conclusion.

2. Interval numbers and their operation

In this section, some interval definitions and interval operation are presented, these preliminaries will be used in the next section [20-22].

Definition 1. Let $a = [a^l, a^u] = \{x \in \square \mid a^l \leq x \leq a^u, a^l, a^u \in \square\}$, then a is called an interval number where a^l, a^u represent the lower and upper bound of a , respectively. If $a^l = a^u$ then a is a real number. Also, if $a^l > 0$ then a is called a positive interval number and if $a^l \geq 0$ then a is called a non-negative interval number.

The set of all interval numbers in $\square$ is represented by $I(\square)$.

Let $a = [a^l, a^u]$ and $b = [b^l, b^u]$ are interval numbers, and $\lambda \geq 0$ is a real number, then:

1. $a = b$ if and only if $a^l = b^l$ and $a^u = b^u$
2. $a + b = [a^l + b^l, a^u + b^u]$
3. $a - b = [a^l - b^u, a^u - b^l]$
4. $\lambda a = \lambda[a^l, a^u] = [\lambda a^l, \lambda a^u]$. Specially $\lambda a = 0$ if $\lambda = 0$
5. $a * b = [\min\{a^l b^l, a^l b^u, a^u b^l, a^u b^u\}, \max\{a^l b^l, a^l b^u, a^u b^l, a^u b^u\}]$
6. $a^2 = [\max\{0, \min\{a^l a^l, a^l a^u, a^u a^u\}\}, \max\{a^l a^l, a^l a^u, a^u a^u\}]$

7. If $a^l \geq 0$, then $\sqrt{a} = [\sqrt{a^l}, \sqrt{a^u}]$.

Definition 2. Let $a_i = [a_i^l, a_i^u]$ ($i = 1, 2, \dots, n$) be interval numbers then $A = \{a_1, a_2, \dots, a_n\}$ is an interval n -dimensional vector.

Definition 3. Let $a_{ij} = [a_{ij}^l, a_{ij}^u]$ ($i = 1, 2, \dots, m; j = 1, 2, \dots, n$) be interval numbers then $A = (a_{ij})_{m \times n}$ is an interval $m \times n$ matrix.

Definition 4. Let $a = [a^l, a^u]$ and $b = [b^l, b^u]$ be interval numbers, $l_a = a^u - a^l$, $l_b = b^u - b^l$ then the degree of possibility of $a \geq b$ is defined as follows:

$$p(a \geq b) = \max \left\{ 1 - \max \left(\frac{b^u - a^l}{l_a + l_b}, 0 \right), 0 \right\} \quad (1)$$

and satisfies following properties:

- a. $0 \leq p(a \geq b) \leq 1$
- b. $p(a \geq b) = 0$ if and only if $a^u \leq b^l$
- c. $p(a \geq b) = 1$ if and only if $b^u \leq a^l$
- d. $p(a \geq a) = \frac{1}{2}$
- e. $p(a \geq b) + p(b \geq a) = 1$

To rank the interval numbers $a_i = [a_i^l, a_i^u]$ ($i = 1, 2, \dots, n$), first the interval numbers a_i ($i = 1, 2, \dots, n$) are compared pairwise using formula (1) and let $p_{it} = p(a_i \geq a_t)$ $i, t = 1, 2, \dots, n$. Then a complementary matrix is constructed as follows:

$$P = (p_{it})_{n \times n} \quad (2)$$

Summing all elements in each row of matrix P , we have:

$$p_i = \sum_{t=1}^n p_{it} \quad i = 1, 2, \dots, n \quad (3)$$

Then the interval numbers a_i ($i = 1, 2, \dots, n$) can be reordered in descending order in accordance with the values of p_i s.

3. The proposed method

In this section the proposed method and its algorithm are presented. Suppose $X = \{x_1, x_2, \dots, x_I\}$ ($I \geq 2$) be a set of alternatives which we want to rank them according to their performance in terms of the set of criteria $C = \{c_1, c_2, \dots, c_J\}$. Let w_j ($j = 1, 2, \dots, J$) be the weight of

the criterion c_j where $w_j \geq 0$ ($j = 1, 2, \dots, J$), $\sum_{j=1}^J w_j = 1$, and

$x_{ij} = [x_{ij}^l, x_{ij}^u]$ ($i = 1, 2, \dots, I, j = 1, 2, \dots, J$) be the performance of i th alternative under j th criterion.

First, to normalize the data, we define:

$$y_{ij} = [y_{ij}^l, y_{ij}^u] \quad i = 1, 2, \dots, I; j = 1, 2, \dots, J \quad (4)$$

where

$$\begin{aligned} y_{ij}^l &= \frac{x_{ij}^l - x_j^-}{x_j^+ - x_j^-} \\ y_{ij}^u &= \frac{x_{ij}^u - x_j^-}{x_j^+ - x_j^-} \end{aligned} \quad j \in O$$

and

$$\begin{aligned} y_{ij}^l &= \frac{x_j^+ - x_{ij}^u}{x_j^+ - x_j^-} \\ y_{ij}^u &= \frac{x_j^+ - x_{ij}^l}{x_j^+ - x_j^-} \end{aligned} \quad j \in O'$$

where O is associated with benefit criteria, these criteria whatever be more, it is better and they are considered as output. Also O' is associated with cost criteria, these criteria whatever be less, it is better and they are considered as input.

and

$$\begin{aligned} x_j^+ &= \max_{i=1, 2, \dots, I} \{x_{ij}^u\} \\ x_j^- &= \min_{i=1, 2, \dots, I} \{x_{ij}^l\} \end{aligned}$$

Then g is denoted the PIS which is an interval vector and obtained as follows:

$$g = (g_1, g_2, \dots, g_J)^T \quad (5)$$

where

$$g_j = \left[\max_{i=1, 2, \dots, I} y_{ij}^l, \max_{i=1, 2, \dots, I} y_{ij}^u \right]. \quad j = 1, 2, \dots, J$$

Now the separation of the alternative x_i ($i = 1, 2, \dots, I$) from the PIS is calculated as follows (using the interval calculations in section 2):

$$d_i^+ = \left[d_i^{+(l)}, d_i^{+(u)} \right] = \sqrt{\sum_{j=1}^J W_j^2 \left[g_j^l - y_{ij}^u, g_j^u - y_{ij}^l \right]^2} \quad i = 1, 2, \dots, I \quad (6)$$

Next we define:

$$d^+ = \min_{i=1,2,\dots,I} \left\{ d_i^{+(l)} \right\} \quad (7)$$

In fact, d^+ shows the least possible separation of an alternative to the PIS.

The NIS is showed by $b = (b_1, b_2, \dots, b_J)^T$, where

$$b_j = \left[\min_{i=1,2,\dots,I} y_{ij}^l, \min_{i=1,2,\dots,I} y_{ij}^u \right] \quad j = 1, 2, \dots, J \quad (8)$$

Similarly the separation of the alternative x_i ($i = 1, 2, \dots, I$) from the NIS is computed as follows (using the interval calculation in section 2):

$$d_i^- = \left[d_i^{-(l)}, d_i^{-(u)} \right] = \sqrt{\sum_{j=1}^J w_j^2 \left[y_{ij}^l - b_j^u, y_{ij}^u - b_j^l \right]^2} \quad i = 1, 2, \dots, I \quad (9)$$

Besides, let

$$d^- = \max_{i=1,2,\dots,I} \left\{ d_i^{-(u)} \right\} \quad (10)$$

In fact, d^- is the maximum separation to the NIS for all alternatives.

Now, the relative agreement index of x_i ($i = 1, 2, \dots, I$) is defined as follows:

$$R_i = \frac{d_i^-}{d^-} - \frac{d_i^+}{d^+} \quad i = 1, 2, \dots, I \quad (11)$$

Actually, R_i is an interval number where:

$$R_i = \left[R_i^l, R_i^u \right] \quad (12)$$

and

$$R_i^l = \frac{d_i^{-(l)}}{d^-} - \frac{d_i^{+(u)}}{d^+} \quad \& \quad R_i^u = \frac{d_i^{-(u)}}{d^-} - \frac{d_i^{+(l)}}{d^+}$$

Remark 1: Generally, the values of d^+ and d^- are positive and only in some special cases they can be zero. In such cases, to obtain R_i ($i = 1, 2, \dots, I$) formula (13) is used instead of formula (12):

$$R_i = \left[R_i^l, R_i^u \right] \quad (13)$$

where

$$R_i^l = \frac{d_i^{-(l)} + 1}{d^- + 1} - \frac{d_i^{+(u)} + 1}{d^+ + 1} \quad \& \quad R_i^u = \frac{d_i^{-(u)} + 1}{d^- + 1} - \frac{d_i^{+(l)} + 1}{d^+ + 1}$$

Actually, with this conversion, the separation of alternatives to the PIS (d_i^+) and to NIS (d_i^-) are summed with the interval number $[1,1]$. Therefore, d^+ and d^- move one unit to right (see Example 3). So, an index called relative agreement index ($R_i; i = 1, 2, \dots, I$) is assigned to each alternative, where $R_i^u \leq 0$. The higher relative agreement of an alternative, means that it is closer to the PIS and farther from the NIS. Thus R_i ($i = 1, 2, \dots, I$) shows the satisfactory level for both criteria of the shortest distance from the PIS and the remotest distance from the NIS.

To compare the alternatives, the interval numbers (R_i s) should be compared (as mentioned in section 2).

For this purpose, the complimentary matrix P is constructed using formula (1) as follows:

$$P = (p_{it})_{I \times I} \quad i, t = 1, 2, \dots, I \quad (14)$$

where

$$p_{it} = p(R_i \geq R_t)$$

then summing up all elements in each row of matrix P , we have:

$$p_i = \sum_{t=1}^I p_{it} \quad i = 1, 2, \dots, I \quad (15)$$

Now, all alternatives x_i ($i = 1, 2, \dots, I$) are ranked according to p_i ($i = 1, 2, \dots, I$) in descending order.

Briefly, the algorithm of the proposed method is summarized by the following steps:

Step 1: Normalize the data using formula (4).

Step 2: Apply formula (5) to determine the PIS (g).

Step 3: Calculate the separation of the alternative x_i ($i = 1, 2, \dots, I$) from the PIS (d_i^+) using formula (6).

Step 4: Obtain d^+ by formula (7).

Step 5: Apply formula (8) to determine the NIS (b).

Step 6: Calculate the separation of the alternative x_i ($i = 1, 2, \dots, I$) from the NIS (d_i^-) using formula (9).

Step 7: Obtain d^- by formula (10).

Step 8: Apply formula (12) to calculate relative agreement index of x_i ($i = 1, 2, \dots, I$) (R_i).

Step 9: Construct complimentary matrix P using formula (14) to compare interval numbers R_i ($i = 1, 2, \dots, I$).

Step 10: Calculate p_i ($i = 1, 2, \dots, I$) using formula (15) and rank all alternatives (x_i s) ($i = 1, 2, \dots, I$) according to p_i ($i = 1, 2, \dots, I$) in descending order.

4. Comparison

Until now, many methods have been presented to solve the interval MCDM problems [14-19]. The main difference between these methods is in normalizing decision matrix and used aggregating functions. The interval TOPSIS method proposed by Jahanshahloo et al. [15] is one of the most popular methods. In this section first, the interval TOPSIS method presented by Jahanshahloo et al. [15] is outlined and then compared by the proposed method. Briefly, the algorithm of the interval TOPSIS method is as follows:

Step 1: Calculate the normalized decision matrix as follows:

$$n_{ij} = [n_{ij}^l, n_{ij}^u] \quad i = 1, 2, \dots, I, \quad j = 1, 2, \dots, J$$

where

$$n_{ij}^l = \frac{x_{ij}^l}{\sqrt{\sum_{i=1}^I [(x_{ij}^l)^2 + (x_{ij}^u)^2]}}$$

$$n_{ij}^u = \frac{x_{ij}^u}{\sqrt{\sum_{i=1}^I [(x_{ij}^l)^2 + (x_{ij}^u)^2]}}$$

Step 2: Construct the weighted normalized decision matrix as $v_{ij}^l = w_j n_{ij}^l$ and $v_{ij}^u = w_j n_{ij}^u$ for $i = 1, 2, \dots, I$

and $j = 1, 2, \dots, J$, where w_j ($j = 1, 2, \dots, J$) is the weight of j th criterion and $\sum_{j=1}^J w_j = 1$.

Step 3: Identify the PIS of k th alternative as follows:

$$A_k^+ = [A_k^{+l}, A_k^{+u}] \quad k = 1, 2, \dots, I$$

where

$$A_k^{+l} = \left\{ (v_1^{+l}, v_2^{+l}, \dots, v_J^{+l}) \right\} = \left\{ \left(\max_{i \neq k} \{v_{ij}^u, v_{kj}^l\} \mid j \in O\}, \left(\min_{i \neq k} \{v_{ij}^l, v_{kj}^u\} \mid j \in O'\} \right) \right\}$$

$$A_k^{+u} = \left\{ (v_1^{+u}, v_2^{+u}, \dots, v_J^{+u}) \right\} = \left\{ \left(\max_{i \neq k} v_{ij}^u \mid j \in O\}, \left(\min_{i \neq k} v_{ij}^l \mid j \in O'\} \right) \right\}$$

Step 4: Identify the NIS of k th alternative as follows:

$$A_k^- = [A_k^{-l}, A_k^{-u}] \quad k = 1, 2, \dots, I$$

where

$$A_k^{-l} = \left\{ (v_1^{-l}, v_2^{-l}, \dots, v_J^{-l}) \right\} = \left\{ \left(\min_{i \neq k} v_{ij}^l \mid j \in O\}, \left(\max_{i \neq k} v_{ij}^u \mid j \in O'\} \right) \right\}$$

$$A_k^{-u} = \left\{ (v_1^{-u}, v_2^{-u}, \dots, v_J^{-u}) \right\} = \left\{ \left(\min_{i \neq k} \{v_{ij}^l, v_{kj}^u\} \mid j \in O\}, \left(\max_{i \neq k} \{v_{ij}^u, v_{kj}^l\} \mid j \in O'\} \right) \right\}$$

Step 5: Calculate the separation of k th alternative from its PIS as:

$$d_k^+ = [d_k^{+l}, d_k^{+u}] \quad k = 1, 2, \dots, I$$

where

$$d_k^{+l} = \left[\sum_{j \in O'} (v_j^{+l} - v_{kj}^l)^2 + \sum_{j \in O} (v_j^{+l} - v_{kj}^u)^2 \right]^{1/2}$$

$$d_k^{+u} = \left[\sum_{j \in O'} (v_j^{+u} - v_{kj}^u)^2 + \sum_{j \in O} (v_j^{+u} - v_{kj}^l)^2 \right]^{1/2}$$

Step 6: Calculate the separation of k th alternative from its NIS as:

$$d_k^- = [d_k^{-l}, d_k^{-u}] \quad k = 1, 2, \dots, I$$

where

$$d_k^{-l} = \left[\sum_{j \in O'} (v_j^{-l} - v_{kj}^l)^2 + \sum_{j \in O} (v_j^{-l} - v_{kj}^u)^2 \right]^{1/2}$$

$$d_k^{-u} = \left[\sum_{j \in O'} (v_j^{-u} - v_{kj}^u)^2 + \sum_{j \in O} (v_j^{-u} - v_{kj}^l)^2 \right]^{1/2}$$

Step 7: Define the relative closeness of k th alternative to its PIS as:

$$R_k = [R_k^l, R_k^u] \quad k = 1, 2, \dots, I$$

where

$$R_k^l = \frac{d_k^{-l}}{d_k^{-u} + d_k^{+u}}$$

$$R_k^u = \frac{d_k^{-u}}{d_k^{-l} + d_k^{+l}}$$

Step 8: Rank the preference order of all the alternatives according to the R_k s ($k = 1, 2, \dots, I$).

To make a comparison between ours and the interval TOPSIS method, the normalization step and the used aggregating function are considered in both methods.

4.1. The effect of Normalization

Generally, in MCDM problems the normalization (scaling) is employed to remove dimensions of criteria values. Now let the original data is transformed as below:

$$x'_{ij} = \alpha x_{ij} + \beta \quad \alpha, \beta \in \mathbb{R} \quad \alpha > 0. \quad (16)$$

If x_{ij} be an interval number, by displaying β in interval form, $[\beta, \beta]$, (as mentioned in section 2) we have:

$$[x'_{ij}{}^l, x'_{ij}{}^u] = [\alpha x_{ij}^l + \beta, \alpha x_{ij}^u + \beta] \quad i = 1, 2, \dots, I, \quad j = 1, 2, \dots, J \quad \beta \in I(\mathbb{R}), \alpha \in \mathbb{R}, \alpha > 0 \quad (17)$$

Now the question is that does this translation affect the decision results obtained by these two methods? In the proposed method, the normalized values of x_{ij} ($i = 1, 2, \dots, I, j = 1, 2, \dots, J$) are calculated using

formula (4). Therefore the normalized criterion values of x'_{ij} ($i=1,2,\dots,I, j=1,2,\dots,J$) are calculated as follows:

$$y'_{ij} = [y'^l_{ij}, y'^u_{ij}] \quad (18)$$

where

$$\begin{aligned} y'^l_{ij} &= \frac{x'^l_{ij} - x^-_j}{x^+_{ij} - x^-_j} \\ y'^u_{ij} &= \frac{x'^u_{ij} - x^-_j}{x^+_{ij} - x^-_j} \end{aligned} \quad j \in O$$

and

$$\begin{aligned} y'^l_{ij} &= \frac{x^+_{ij} - x'^u_{ij}}{x^+_{ij} - x^-_j} \\ y'^u_{ij} &= \frac{x^+_{ij} - x'^l_{ij}}{x^+_{ij} - x^-_j} \end{aligned} \quad j \in O'$$

where

$$\begin{aligned} x^+_{ij} &= \max_{i=1,2,\dots,I} \{x'^u_{ij}\} \\ x^-_{ij} &= \min_{i=1,2,\dots,I} \{x'^l_{ij}\} \end{aligned}$$

By formula (17) we have:

$$\begin{aligned} x^+_{ij} &= \max_{i=1,2,\dots,I} \{\alpha x'^u_{ij} + \beta\} = \alpha \max_{i=1,2,\dots,I} \{x'^u_{ij}\} + \beta = \alpha x^+_{ij} + \beta \\ x^-_{ij} &= \min_{i=1,2,\dots,I} \{\alpha x'^l_{ij} + \beta\} = \alpha \min_{i=1,2,\dots,I} \{x'^l_{ij}\} + \beta = \alpha x^-_{ij} + \beta \end{aligned} \quad (19)$$

Now using formulas (17) and (19) we have:

$$\begin{aligned} y'^l_{ij} &= \frac{\alpha x'^l_{ij} + \beta - \alpha x^-_j - \beta}{\alpha x^+_{ij} + \beta - \alpha x^-_j - \beta} = \frac{x'^l_{ij} - x^-_j}{x^+_{ij} - x^-_j} = y'^l_{ij} \\ y'^u_{ij} &= \frac{\alpha x'^u_{ij} + \beta - \alpha x^-_j - \beta}{\alpha x^+_{ij} + \beta - \alpha x^-_j - \beta} = \frac{x'^u_{ij} - x^-_j}{x^+_{ij} - x^-_j} = y'^u_{ij} \end{aligned} \quad j \in O \quad i = 1, 2, \dots, I \quad (20)$$

A similar discussion is hold for negative criteria ($j \in O'$) as well. Thus:

$$y'_{ij} = y_{ij} \quad i = 1, 2, \dots, I; j = 1, 2, \dots, J \quad (21)$$

This shows that the proposed method is translation invariance. That is, the normalized criteria values in our method is independent of the measurement units and dimensions of criteria.

In the proposed method by Jahanshahloo et al. [15], the normalized criteria values of x_{ij} ($i = 1, 2, \dots, I, j = 1, 2, \dots, J$) are calculated as:

$$n_{ij} = [n_{ij}^l, n_{ij}^u] \quad i = 1, 2, \dots, I; j = 1, 2, \dots, J \quad (22)$$

where

$$n_{ij}^l = \frac{x_{ij}^l}{\sqrt{\sum_{i=1}^I [(x_{ij}^l)^2 + (x_{ij}^u)^2]}}$$

$$n_{ij}^u = \frac{x_{ij}^u}{\sqrt{\sum_{i=1}^I [(x_{ij}^l)^2 + (x_{ij}^u)^2]}}$$

Hence the normalized criteria values of $x_{ij}^{'}$ ($i = 1, 2, \dots, I, j = 1, 2, \dots, J$) are calculated as follows:

$$n_{ij}^{' } = [n_{ij}^{'l}, n_{ij}^{'u}] \quad i = 1, 2, \dots, I, \quad j = 1, 2, \dots, J \quad (23)$$

where

$$n_{ij}^{'l} = \frac{x_{ij}^{'l}}{\sqrt{\sum_{i=1}^I [(x_{ij}^{'l})^2 + (x_{ij}^{'u})^2]}}$$

$$n_{ij}^{'u} = \frac{x_{ij}^{'u}}{\sqrt{\sum_{i=1}^I [(x_{ij}^{'l})^2 + (x_{ij}^{'u})^2]}}$$

By formulas (17) and (23) we have:

$$n_{ij}^{'l} = \frac{\alpha x_{ij}^l + \beta}{\sqrt{\sum_{i=1}^I [(\alpha x_{ij}^l + \beta)^2 + (\alpha x_{ij}^u + \beta)^2]}}$$

$$n_{ij}^{'u} = \frac{\alpha x_{ij}^u + \beta}{\sqrt{\sum_{i=1}^I [(\alpha x_{ij}^l + \beta)^2 + (\alpha x_{ij}^u + \beta)^2]}}$$

$$i = 1, 2, \dots, I; j = 1, 2, \dots, J \quad (24)$$

So $n_{ij}^{' } \neq n_{ij}$ if $\beta \neq [0, 0]$, i.e. the normalized criterion values in the interval TOPSIS method could depend on the evaluation units and dimensions of criterion values.

4.2. Aggregating function

The proposed method employs the aggregating function as formula (12), which demonstrates the extent that each alternative is close to the PIS and is remote from the NIS. In fact, the selected alternative has the maximum value of R_i ($i = 1, 2, \dots, I$). Evidently, this alternative would like to minimize its distance from

the PIS (d_i^+ , $i = 1, 2, \dots, I$) and maximize the distance from the NIS (d_i^- , $i = 1, 2, \dots, I$) at the same time. In other words, R_i ($i = 1, 2, \dots, I$) indicates the satisfactory degree which the DM considers the compromise extent between the shortest distance from the PIS and the farthest distance from the NIS.

Thus, in the proposed method, the distances of each alternative from the PIS (d_i^+ , $i = 1, 2, \dots, I$) and NIS (d_i^- , $i = 1, 2, \dots, I$) have the same effect in the final ranking, but the interval TOPSIS method is not so. The interval TOPSIS method [15] is based on following aggregating function:

$$R_k = [R_k^l, R_k^u] \quad k = 1, 2, \dots, I \quad (25)$$

where

$$R_k^l = \frac{d_k^{-l}}{d_k^{-u} + d_k^{+u}}$$

$$R_k^u = \frac{d_k^{-u}}{d_k^{-l} + d_k^{+l}}$$

Now, consider k th and j th alternatives. Suppose R_j and R_k are obtained from formula (25) and besides,

$R_j^l < R_k^l$ and $R_j^u < R_k^u$, that means:

$$\frac{d_j^{-l}}{d_j^{-u} + d_j^{+u}} < \frac{d_k^{-l}}{d_k^{-u} + d_k^{+u}} \quad (26)$$

$$\frac{d_j^{-u}}{d_j^{-l} + d_j^{+l}} < \frac{d_k^{-u}}{d_k^{-l} + d_k^{+l}}$$

So using the interval TOPSIS method, k th alternative is ranked higher than j th alternative. In a special case, the inequalities (26) may be satisfied, while $d_j^{+l} < d_k^{+l}$ and $d_j^{+u} < d_k^{+u}$, i.e. the distance of j th alternative to the PIS may be shorter than the distance of k th alternative to the PIS, while k th alternative is higher than j th alternative in ranking (see Example 1). This illogical ranking is due to this fact that the interval TOPSIS method pay more attention to the distance of alternative to the NIS (d_i^- , $i = 1, 2, \dots, I$) than the distance of alternative to the PIS (d_i^+ , $i = 1, 2, \dots, I$). Indeed, this is the disadvantage of the interval TOPSIS method.

5. Illustration

In this section, three examples are considered to illustrate the proposed method and compare our results with those published in literature.

Example 1. The example adopted from Jahanshahloo et al. [15] is considered here. This example is related to the assessment of six cities in Iran to find the best place for creating a data factory. These cities must be evaluated by four criteria which are: Distance from border (C_1), cost of creating the factory (C_2), Finance

(C_3), and Product in the region (C_4). C_1 and C_2 are cost criteria, C_3 and C_4 are benefit criteria. Table 1 represents the data. The first criterion is a real number and the others are in interval form and all the criteria have the same importance i.e. $w_j = 0.25 (j = 1, 2, 3, 4)$.

Using the proposed method in step 1 data are normalized by formula (4) (Table 2). In steps 2, 3 the PIS (g) is determined by formula (5) (Table 3) then $d_i^+ (i = 1, 2, \dots, 6)$ is calculated for all alternatives using formula (6) (Table 4). In step 4 by formula (7), we have $d^+ = 0.1273$. In steps 5, 6 the NIS (b) is determined using formula (8) (Table 3), then $d_i^- (i = 1, 2, \dots, 6)$ is calculated for all alternatives using formula (9) (Table 4). In step 7 by formula (10) we have $d^- = 0.3984$. By formula (12) in step 8, $R_i (i = 1, 2, \dots, 6)$ values are calculated (Table 5). In the next steps, to compare interval numbers $R_i (i = 1, 2, \dots, 6)$ and rank the alternatives, complementary matrix P is constructed using formula (14), then p_i values ($i = 1, 2, \dots, 6$) are calculated using formula (15) (shown in Table 5) and all alternatives are ranked according to $p_i (i = 1, 2, \dots, 6)$ in descending order.

Ranking obtained from the proposed method and Jahanshahloo et al. [15], are shown in Table 6. As seen in Table 4, the distance of the second alternative to the PIS is shorter than those of the fifth alternative, but the proposed method in the mentioned paper puts the fifth alternative in the third rank and the second alternative in the fourth rank. Also, the distance of the sixth alternative to the PIS is shorter than those of the fourth alternative, but the method proposed by Jahanshahloo et al. puts the fourth alternative in the first rank and the sixth one in the second rank. The reason for this inappropriate ranking by Jahanshahloo et al. is the same as mentioned before. That is, paying more attention to distance to the NIS than the distance to the PIS. This drawback leads to the situation that an alternative with shorter distance to the PIS, ranked in lower rank. But as seen in Table 6, the proposed method ranks the second alternative higher than the fifth one and the sixth alternative is ranked higher than the fourth one which this ranking is more logical. Meanwhile, in the method proposed by Jahanshahloo et al., the PIS and the NIS change for each alternative, so that the relationship between the alternatives is gone and the final comparison will not be relative.

Example 2. Here, the example provided by Sayadi et al. [16] is considered. In this example there are three alternatives and two criteria. C_1 is a cost criterion and C_2 is a benefit criterion, both criteria have similar relative importance, i.e. $w_j = 0.5 (j = 1, 2)$. The interval decision matrix is shown in Table 7.

To solve this problem using the proposed method, in step 1 data are normalized by formula (4) (Table 8). In steps 2, 3 the PIS (g) is determined using formula (5) (Table 9), then $d_i^+ (i = 1, 2, 3)$ are calculated for all alternatives using formula (6) (Table 10). In step 4 by formula (7) we have $d^+ = 0.1571$. In steps 5, 6

the NIS (b) is determined using formula (8) (Table 9) then $d_i^-(i=1,2,3)$ are calculated for all alternatives by formula (9) (Table 10). In step 7 by formula (10), we have $d^- = 0.5114$. Employing formula (12) in step 8, R_i values ($i=1,2,3$) are calculated (Table 11).

In the next steps, to compare interval numbers $R_i(i=1,2,3)$ and rank the alternatives, complimentary matrix P is constructed by formula (14), then $p_i(i=1,2,3)$ are calculated using formula (15) (shown in Table 11) and all alternatives are ranked according to $p_i(i=1,2,3)$ in descending order. Table 12 shows the ranking obtained from the proposed method and the proposed method by Sayadi et al. [16]. It is observed that the ranking obtained from both method is the same. It should be noted that in the method proposed by Sayadi et al., parameter ν is very influential in the final ranking and its optimal value is not known, moreover ν is considered 0.5 without any justification. Also, indicator α , to compare interval numbers, could significantly affect the final ranking and DM may not be able to determine this value. In fact, it is very difficult to determine both parameters α and ν in real-life decision situations.

Example 3. The example adopted from Dymova et al. [17] is considered. This example deals with three alternatives and four criteria presented by interval numbers in Table 13, where C_1 and C_2 are benefit criteria, C_3 and C_4 are cost criteria. All the criteria have the same importance, i.e. $w_j = 0.25(j=1,2,3,4)$.

Using the proposed method, steps 1 to 3 are executed like previous examples. Obtained values are listed in Tables 14 to 16, respectively. In step 4 we have $d^+ = 0$. Also obtained values in steps 5, 6 are listed in Tables 15, 16. In step 7 we have $d^- = 0.4747$. This example is different from the previous ones in step 8, because in step 4, $d^+ = 0$. Thus, according to Remark 1, formula (12) should be employed instead of (13), to calculate $R_i(i=1,2,3)$, obtained values were shown in Table 17. In the next steps, $p_i(i=1,2,3)$ are calculated (Table 17).

Then all alternatives are ranked in descending order $p_i(i=1,2,3)$. The obtained ranking from both methods are shown in Table 18. As shown, the ranking of the two methods are the same. But in the method proposed by Dymova et al., the normalized criterion values depend on the evaluation units and dimensions of criterion values and to determine the PIS and the NIS and calculate the intervals distance, only the centers of the intervals are taken into account which may result in partial data loss.

6. Conclusion

Due to this fact that in some cases, determining the exact values of the criteria is difficult, it is more appropriate to consider them as interval numbers in MCDM problems. In this paper, a method is presented

to solve MCDM problems with interval numbers. The proposed method provides a balance between closeness to the PIS and being far away from the NIS. Putting in other words, these two attributes have the same effect in alternative ranking. As proved in sub-section 4.1, the proposed method is translation invariance. Moreover, in this method all calculations are done in interval form unlike the papers in which the computations are not completely based on interval arithmetic.

In the proposed method, the PIS and the NIS are determined as interval numbers and calculated the distance of each alternative to the PIS and the NIS by interval extension of the Euclidean norm. Then the relative agreement of each alternative is obtained. These relative agreements are compared together to rank the alternatives. Numerical examples show the advantages of the proposed method compared to other published studies. For future researches, it is suggested that employing interval extension of p -norm instead of interval extension of Euclidean norm to obtain the distances of alternatives to the PIS and the NIS. Also, one can use other methods instead of constructing the complimentary matrix P to compare and sort the interval numbers.

References

- [1] Saaty, T. L. “*The analytical hierarchy process*”, McGraw-Hill, New York (1980).
- [2] Hwang, C. L., & Yoon, K. “Multiple attribute decision making: methods and applications”, *Springer*, New York (1981).
- [3] Zavadskas, E. K., Kaklauskas, A., & Sarka, V. “The new method of multi-criteria complex proportional assessment of projects”, *Technological and economic development of economy*, 1(3), pp. 131-139 (1994).
- [4] Zavadskas, E. K., Antucheviciene, J., Turskis, Z., et al. “Hybrid multiple-criteria decision-making methods: A review of applications in engineering”, *Scientia Iranica* 23(1), pp. 1-20 (2016).
- [5] Govindan, K., & Jepsen, M. B. “ELECTRE: A comprehensive literature review on methodologies and applications”, *European Journal of Operational Research*, 250(1), pp. 1-29 (2016).
- [6] Carneiro, J., Conceição, L., Martinho, D. et al. “Including cognitive aspects in multiple criteria decision analysis”, *Annals of Operations Research*, 265(2), pp. 269-291 (2018).
- [7] MacCrimmon, K. R. “Decision making among multiple-attribute alternatives: a survey and consolidated approach”, *Rand Corp Santa Monica Ca*, (1968).
- [8] Opricovic, S. “Multicriteria optimization of civil engineering systems”, *Faculty of Civil Engineering, Belgrade*, 2(1), pp. 5-21 (1998).
- [9] Abdollahi, A., Pour-Moallem, N., & Abdollahi A. “Dynamic Negawatt Demand Response Resource Modeling and Prioritizing in Power Markets”, *Scientia Iranica*, DOI: 10.24200/sci.2017.4406 (2017).
- [10] Kirubakaran, B., & Ilankumaran, M. “Selection of optimum maintenance strategy based on FAHP integrated with GRA–TOPSIS”, *Annals of Operations Research*, 245(1-2), pp. 285-313 (2016).
- [11] Singh, R. K., Gunasekaran, A., & Kumar, P. “Third party logistics (3PL) selection for cold chain management: a fuzzy AHP and fuzzy TOPSIS approach”, *Annals of Operations Research*, pp. 1-23 (2017).
- [12] Mahmoudi, A., Sadi-Nezhad, S., & Makui, A. “An extended fuzzy VIKOR for group decision-making based on fuzzy distance to supplier selection”, *Scientia Iranica. Transaction E, Industrial Engineering*, 23(4), pp. 1879 (2016).

- [13] Hu, J., Chen, P., & Chen, X. "Intuitionistic random multi-criteria decision-making approach based on prospect theory with multiple reference intervals", *Scientia Iranica. Transaction E, Industrial Engineering*, 21(6), pp. 2347 (2014).
- [14] Jahanshahloo, G. R., Lotfi, F. H., & Izadikhah, M. "An algorithmic method to extend TOPSIS for decision-making problems with interval data", *Applied mathematics and computation*, 175(2), pp. 1375-1384 (2006).
- [15] Jahanshahloo, G. R., Lotfi, F. H., & Davoodi, A. R. "Extension of TOPSIS for decision-making problems with interval data: Interval efficiency", *Mathematical and Computer Modelling*, 49(5-6), pp. 1137-1142 (2009).
- [16] Sayadi, M. K., Heydari, M., & Shahanaghi, K. "Extension of VIKOR method for decision making problem with interval numbers", *Applied Mathematical Modelling*, 33(5), pp. 2257-2262 (2009).
- [17] Dymova, L., Sevastjanov, P., & Tikhonenko, A. "A direct interval extension of TOPSIS method", *Expert Systems with Applications*, 40(12), pp. 4841-4847 (2013).
- [18] Hafezalkotob, A., Hafezalkotob, A., & Sayadi, M. K. "Extension of MULTIMOORA method with interval numbers: an application in materials selection", *Applied Mathematical Modelling*, 40(2), pp. 1372-1386 (2016).
- [19] Jahan, A., & Edwards, K. L. "VIKOR method for material selection problems with interval numbers and target-based criteria", *Materials & Design*, 47, pp. 759-765 (2013).
- [20] Xu, Z. "On method for uncertain multiple attribute decision making problems with uncertain multiplicative preference information on alternatives", *Fuzzy Optimization and Decision Making*, 4(2), pp. 131-139 (2005).
- [21] Xu, Z. "Dependent uncertain ordered weighted aggregation operators" *Information Fusion*, 9(2), pp. 310-316 (2008).
- [22] Saffarzadeh, S., Hadi-Vencheh, A., & Jamshidi, A. "Weight determination and ranking priority in interval group MCDM", *Scientia Iranica* 27(6), pp. 3242-3252 (2020).

Biographies

Saghi Saffarzadeh is an assistant professor of Operations Research. Her fields of study are: multiple criteria decision making, interval programming, fuzzy mathematical programming and data envelopment analysis. She has published some papers in international journals.

Ali Jamshidi is an assistant professor of Operations Research. His research interests include multiple criteria decision modeling, data envelopment analysis, supply chain management and scheduling. He has published several papers in international journals.

Abdollah Hadi-Vencheh is a full professor of Operations Research and Decision Sciences at IAU, Isfahan Branch. His research interests lie in the broad area of multiple criteria decision making, performance management, data envelopment analysis, fuzzy mathematical programming, and fuzzy decision making. He has published more than 100 papers in more prestigious international journals such as *European Journal of Operational Research*, *IEEE Transaction on Fuzzy Systems*, *Information Sciences*, *Computers and Industrial Engineering*, *Journal of the Operational Research Society*, *Journal of manufacturing systems*, *Expert Systems with Applications*, *Expert Systems*, *Measurement*, *Kybernetes*, *Optimization*, *Optimization Letters*, *Scientia Iranica*, *Computers in Industry*, and *International Journal of Computer Integrated Manufacturing*.

Table 1. The interval decision matrix for Example 1

Alternative	C ₁	C ₂	C ₃	C ₄
A ₁	1451	[2551, 3118]	[40, 50]	[153, 187]
A ₂	843	[3742, 4573]	[63, 77]	[459, 561]
A ₃	1125	[3312, 4049]	[48, 58]	[153, 187]
A ₄	55	[5309, 6488]	[72, 88]	[347, 426]
A ₅	356	[3709, 4534]	[59, 71]	[151, 189]
A ₆	391	[4884, 5969]	[72, 88]	[388, 474]

Table 2. The interval normalized decision matrix for Example 1.

Alternative	C ₁	C ₂	C ₃	C ₄
A ₁	0.0000	[0.8560, 1.0000]	[0.0000, 0.2083]	[0.0049, 0.0878]
A ₂	0.4355	[0.4864, 0.6975]	[0.4792, 0.7708]	[0.7512, 1.0000]
A ₃	0.2335	[0.6195, 0.8067]	[0.1667, 0.3750]	[0.0049, 0.0878]
A ₄	1.0000	[0.0000, 0.2995]	[0.6667, 1.0000]	[0.4780, 0.6707]
A ₅	0.7844	[0.4963, 0.7059]	[0.3958, 0.6458]	[0.0000, 0.0927]
A ₆	0.7593	[0.1318, 0.4074]	[0.6667, 1.0000]	[0.5780, 0.7878]

Table 3. The positive ideal solution (PIS) and negative ideal solution (NIS) for Example 1.

Ideal	C ₁	C ₂	C ₃	C ₄
PIS (g)	1.0000	[0.8560, 1.0000]	[0.6667, 1.0000]	[0.7512, 1.0000]
NIS (b)	0.0000	[0.0000, 0.2995]	[0.0000, 0.2083]	[0.0000, 0.0878]

Table 4. Distance of each alternative from the positive ideal solution (PIS) and negative ideal solution (NIS) for Example 1.

Alternative	d_i^+	d_i^-
A ₁	[0.3211, 0.4338]	[0.1391, 0.2563]
A ₂	[0.1466, 0.2392]	[0.2148, 0.3767]
A ₃	[0.2640, 0.3887]	[0.0990, 0.2310]
A ₄	[0.1406, 0.2941]	[0.2918, 0.3984]
A ₅	[0.1773, 0.3226]	[0.2075, 0.3102]
A ₆	[0.1273, 0.2623]	[0.2533, 0.3843]

Table 5. The relative agreement index and P_i of each alternative for Example 1.

Alternative	R_i	P_i
A_1	[-3.0595, -1.8801]	0.9235
A_2	[-1.3405, -0.2062]	4.2226
A_3	[-2.8054, -1.4946]	1.3501
A_4	[-1.5782, -0.1046]	4.0697
A_5	[-2.0140, -0.6149]	3.1687
A_6	[-1.4251, -0.0354]	4.2654

Table 6. Ranking of alternatively in Example 1.

Alternative	Proposed method	Jahanshahloo et al. method
A_1	6	6
A_2	2	4
A_3	5	5
A_4	3	1
A_5	4	3
A_6	1	2

Table 7. The interval decision matrix for Example 2.

Alternative	C_1	C_2
A_1	[0.75, 1.24]	[2784, 3192]
A_2	[1.83, 2.11]	[3671, 3857]
A_3	[4.90, 5.73]	[4409, 4681]

Table 8. The interval normalized decision matrix for Example 2.

Alternative	C_1	C_2
A_1	[0.9016, 1.0000]	[0.0000, 0.2151]
A_2	[0.7269, 0.7831]	[0.4676, 0.5656]
A_3	[0.0000, 0.1667]	[0.8566, 1.0000]

Table 9. The positive ideal solution (PIS) and negative ideal solution (NIS) for Example 2.

Ideal	C_1	C_2
PIS (g)	[0.9016, 1.0000]	[0.8566, 1.0000]
NIS (b)	[0.0000, 0.1667]	[0.0000, 0.2151]

Table 10. Distance of each alternative from the positive ideal solution (PIS) and negative ideal solution (NIS) for Example 2.

Alternative	d_i^+	d_i^-
A ₁	[0.3208, 0.5024]	[0.3675, 0.5114]
A ₂	[0.1571, 0.2992]	[0.3073, 0.4830]
A ₃	[0.3675, 0.5051]	[0.3208, 0.5069]

Table 11. The relative agreement index and P_i of each alternative for Example 2.

Alternative	R_i	P_i
A ₁	[-2.4798, -1.0420]	1.1749
A ₂	[-1.3038, -0.0556]	2.4025
A ₃	[-2.5882, -1.3481]	0.9226

Table 12. Ranking of alternatively in Example 2.

Alternative	Proposed method	Sayadi et al. method
A ₁	2	2
A ₂	1	1
A ₃	3	3

Table 13. The interval decision matrix for Example 3.

Alternative	C ₁	C ₂	C ₃	C ₄
A ₁	[6, 22]	[10, 15]	[16, 21]	[18, 20]
A ₂	[15, 18]	[8, 11]	[20, 30]	[19, 28]
A ₃	[9, 13]	[12, 17]	[42, 48]	[40, 49]

Table 14. The interval normalized decision matrix for Example 3.

Alternative	C ₁	C ₂	C ₃	C ₄
A ₁	[0.0000, 1.0000]	[0.2222, 0.7778]	[0.8438, 1.0000]	[0.9355, 1.0000]
A ₂	[0.5625, 0.7500]	[0.0000, 0.3333]	[0.5625, 0.8750]	[0.6774, 0.9677]
A ₃	[0.1875, 0.4375]	[0.4444, 1.0000]	[0.0000, 0.1875]	[0.0000, 0.2903]

Table 15. The positive ideal solution (PIS) and negative ideal solution (NIS) for Example 3.

PIS (g)	[0.5625, 1.0000]	[0.4444, 1.0000]	[0.8438, 1.0000]	[0.9355, 1.0000]
NIS (b)	[0.0000, 0.4375]	[0.0000, 0.3333]	[0.0000, 0.1875]	[0.0000, 0.2903]

Table 16. Distance of each alternative from the positive ideal solution (PIS) and negative ideal solution (NIS) for Example 3.

Alternative	d_i^+	d_i^-
A ₁	[0.0000, 0.3195]	[0.2301, 0.4747]
A ₂	[0.0278, 0.3048]	[0.1383, 0.3853]
A ₃	[0.2322, 0.4308]	[0.0278, 0.2862]

Table 17. The relative agreement index and P_i of each alternative for Example 3.

Alternative	R_i	P_i
A ₁	[-0.4854, 0.0000]	1.9271
A ₂	[-0.5329, -0.0884]	1.7156
A ₃	[-0.7338, -0.3600]	0.8573

Table 18. Ranking of alternatively in Example 3.

Alternative	Proposed method	Dymova et al. method
A ₁	1	1
A ₂	2	2
A ₃	3	3