



Research Note

An efficient synthesis of monoarylidene derivatives of pyran-4-one and piperidin-4-one

M.M. Mojtahedi*, M.S. Abaee, M. Samianifard and A. Shamloo

Chemistry and Chemical Engineering Research Center of Iran, Tehran, P.O. Box 14335-186, Iran.

Received 9 January 2013; received in revised form 22 July 2013; accepted 22 February 2014

KEYWORDS

α, β -unsaturated ketones;
 Organocatalysis;
 Heterocyclic chemistry;
 Aldol condensation.

Abstract. A room-temperature procedure is developed for the direct synthesis of monoarylidene derivatives of pyran-4-one and piperidin-4-one systems under solvent-free conditions. Relatively high yields of products are obtained from the reaction of ketones **1** with different aromatic and aliphatic aldehydes under the catalytic system of TMSNMe₂ and MgBr₂.OEt₂. Notably, formation of the undesired bis counterparts, as a major limitation of these reaction types, is minimized using the employed conditions.

© 2014 Sharif University of Technology. All rights reserved.

1. Introduction

α, β -Unsaturated derivatives of ketones are very useful species in synthetic organic chemistry since they are the key substructure of many natural and biologically active compounds [1-4]. Moreover, these derivatives could be employed as useful precursors in other synthetic transformations [5-6]. Although various methods are available for the synthesis of these compounds [7-9], many of them involve multi-step procedures or demand the use of reactants or reagents which are not commercially available.


Claisen-Schmidt condensation is often a common pathway for the synthesis of α, β -unsaturated carbonyl systems via the reaction of aromatic aldehydes with ketones at their α position [10-14]. However, this approach requires the use of strong bases or acids and is often accompanied by competitive formation of the respective bis by-products. To overcome these

drawbacks, recently, two independent groups reported new procedures catalyzed by *N, N*-dimethylammonium *N', N'*-dimethylcarbamate [15] and a pyrrolidine imide [16], respectively. These methods are based on catalytic cycles with proposed iminium transients for the conversion of reactants to the desired products.

In continuation of our investigations into aldol condensation reactions of various homo- and heterocyclic ketones [17-19], we have decided to apply our experience to the synthesis of the monoarylidene derivatives of such ketones [20,21]. On this basis, in this paper, we have reported the results achieved in pyran-4-one and piperidin-4-one heterocyclic systems (Scheme 1).

It is noteworthy that derivatives of the piperidinone system are considered as important six-membered nitrogen-containing heterocycles, due to possessing diverse biological properties, and which are widely found in natural resources [22-24]. The same case is applied to the pyran ring [25-28]. Consequently, new routes to access derivatives of these systems are nowadays of prominence in synthetic organic chemistry.

*. Corresponding author. Tel.: +98 21 44580749;
 Fax: +98 21 44580785
 E-mail address: mojtahedi@ccerci.ac.ir (M.M. Mojtahedi)



Scheme 1. General reaction for the synthesis of products **3-4**.

2. Experimental

2.1. General

Melting points are uncorrected. IR spectra were recorded using KBr disks on a Bruker Vector-22 infrared spectrometer. NMR spectra were obtained on a Bruker Ultra Shield™ (500 MHz) as CDCl₃ solutions, using TMS as an internal standard reference. Elemental analyses were performed using a Thermo Finnigan Flash EA 1112 instrument. MS spectra were obtained on a Fisons 8000 Trio instrument at ionization potential of 70 eV. TLC experiments were carried out on pre-coated silica gel plates using petroleum ether/EtOAc (4:1) as the eluent. A known procedure was used for the preparation of MgBr₂OEt₂ [29]. Solvents, TMSNMe₂, and other starting materials were purchased from commercial sources. Aldehydes were redistilled or recrystallized before being used. Products **3a** [20] are reported previously. All other products are new and their structures are determined based on their physical and spectroscopic specifications, as listed below.

2.2. Typical procedure

A mixture of an aldehyde (3.0 mmol), ketone **1** (3.0 mmol), MgBr₂OEt₂ (7 mol%, 54 mg), and TMSNMe₂ (176 mg, 237 μ l, 1.5 mmol) was stirred at room temperature under an inert atmosphere for a given period of time. The progress of the reaction was monitored by TLC experiments. At the end, water (5 mL) was added to the mixture and the product was extracted with Et₂O (2 \times 5 mL). The organic layer was dried over a column of Na₂SO₄. Evaporation of the solvent led to a residue, which was purified by column chromatography using silica gel and petroleum ether/EtOAc (4:1) as the eluent.

2.3. Spectral data of new products

(E)-3-(2-Fluorobenzylidene)dihydro-2H-pyran-4(3H)-one (3b). White solid was obtained in 75% yield. Mp 92–93°C; ¹H NMR (500 MHz, CDCl₃) δ 2.68–2.70 (m, 2H), 4.06–4.09 (m, 2H), 4.70 (dd, J = 1.0, 1.5 Hz, 2H), 7.08–7.17 (m, 3H), 7.33–7.36 (m, 1H), 7.66 (d, J = 1.5 Hz, 1H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 40.4, 66.1, 69.0 (d, J = 4.0 Hz), 116.4 (d, J = 22.0 Hz), 122.7 (d, J = 13.5 Hz), 124.5 (d, J = 4.0 Hz), 128.8 (d, J = 4.0 Hz), 131.1 (d, J = 2.0 Hz), 131.7 (d, J = 8.5 Hz), 135.6, 161.3 (d, J = 250.0 Hz), 196.3 ppm; IR (neat) ν 1691, 1618, 1220 cm^{−1}; MS (70 eV): m/z 206 (M⁺), 178, 149, 133. Anal.

Calcd for C₁₂H₁₁FO₂: C, 69.89, H, 5.38. Found: C 69.57; H, 5.46.

(E)-3-(4-Chlorobenzylidene)dihydro-2H-pyran-4(3H)-one (3c). White solid was obtained in 70% yield. Mp 106–107°C; ¹H NMR (500 MHz, CDCl₃) δ 2.67–2.69 (m, 2H), 4.06–4.08 (m, 2H), 4.81 (s, 2H), 7.12 (d, J = 8.5 Hz, 2H), 7.37 (d, J = 8.5 Hz, 2H), 7.55 (s, 1H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 40.2, 65.9, 69.0, 129.4, 132.2, 133.2, 134.1, 135.1, 136.0, 196.3 ppm; IR (neat) ν 2861, 1683, 1596, 1235 cm^{−1}; MS (70 eV): m/z 222 (M⁺), 166, 131, 115. Anal. Calcd for C₁₂H₁₁ClO₂: C, 64.73; H, 4.98. Found: C, 64.34; H, 4.97.

(E)-Methyl 4-((4-oxodihydro-2H-pyran-3(4H)-ylidene)methyl)benzoate (3d). Yellow solid was obtained in 73% yield. Mp 121–122°C; ¹H NMR (500 MHz, CDCl₃) δ 2.69–2.71 (m, 2H), 3.92 (s, 3H), 4.07–4.09 (m, 2H), 4.82 (d, J = 2.0 Hz, 2H), 7.32 (d, J = 8.5 Hz, 2H), 7.60 (d, J = Hz, 1H), 8.05 (d, J = 8.5 Hz, 2H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 40.2, 52.7, 65.9, 69.0, 130.2, 130.7, 130.9, 134.9, 135.4, 139.0, 166.8, 196.3 ppm; IR (neat) ν 1726, 1688, 1284 cm^{−1}; MS (70 eV): m/z 246 (M⁺), 231, 187, 175, 131. Anal. Calcd for C₁₄H₁₄O₄: C, 68.28, H, 5.73. Found: C 68.15; H, 5.78.

(E)-3-(4-Nitrobenzylidene)dihydro-2H-pyran-4(3H)-one (3e). Yellow solid was obtained in 76% yield. Mp 135–136°C; ¹H NMR (500 MHz, CDCl₃) δ 2.71–2.74 (m, 2H), 4.09–4.12 (m, 2H), 4.82 (d, J = 2.0 Hz, 2H), 7.41 (d, J = 8.5 Hz, 2H), 7.60 (d, J = 2 Hz, 1H), 8.26 (d, J = 8.5 Hz, 2H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 40.3, 65.9, 68.8, 124.3, 131.3, 133.3, 136.8, 141.0, 146.0, 196.0 ppm; IR (neat) ν 1687, 1509, 1347 cm^{−1}; MS (70 eV): m/z 233 (M⁺), 216, 205, 160, 130. Anal. Calcd for C₁₂H₁₁NO₄: C, 61.80; H, 4.75. Found: C 61.96; H, 4.85.

(E)-3-(2-Methoxybenzylidene)dihydro-2H-pyran-4(3H)-one (3f). Yellow solid was obtained in 68% yield. Mp 102–103°C; ¹H NMR (500 MHz, CDCl₃) δ 2.67–2.69 (m, 2H), 3.85 (s, 3H), 4.06–4.09 (m, 2H), 4.73 (d, J = 2.0 Hz, 2H), 6.92 (d, J = 8.0 Hz, 1H), 6.96 (d, J = 7.5 Hz, 1H), 7.02 (dd, J = 1.5, 7.5 Hz, 1H), 7.35 (ddd, J = 1.5, 7.5, 8.0 Hz, 1H), 7.85 (m, 1H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 40.4, 55.9, 66.2, 69.2, 111.2, 120.5, 123.7, 130.9, 131.5, 132.2, 133.6, 158.9, 196.9 ppm; IR (neat) ν 2948, 1689, 1459, 1255 cm^{−1}; MS (70 eV): m/z 218 (M⁺), 187, 131, 119. Anal. Calcd for C₁₃H₁₄O₃: C, 71.54, H, 6.47. Found: C 71.11; H, 6.50.

(E)-3-(4-Methoxybenzylidene)dihydro-2H-pyran-4(3H)-one (3g). Yellow solid was obtained in

86% yield. Mp 75–76°C; ^1H NMR (500 MHz, CDCl_3) δ 2.65–2.67 (m, 2H), 3.83 (s, 3H), 4.05–4.07 (m, 2H), 4.85 (d, $J = 2.0$ Hz, 2H), 6.92 (d, $J = 8.6$ Hz, 2H), 7.25 (d, $J = 8.6$ Hz, 2H), 7.59 (d, $J = 2.0$ Hz, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 40.0, 55.8, 65.8, 69.3, 114.7, 127.5, 131.5, 133.1, 136.4, 161.1, 196.3 ppm; IR (neat) ν 1681, 1591, 1511 cm^{-1} ; MS (70 eV): m/z 218 (M^+), 187, 162, 131, 119. Anal. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_3$: C, 71.54, H, 6.47. Found: C 71.21; H, 6.55.

(E)-3-(Furan-2-ylmethylene)dihydro-2H-pyran-4(3H)-one (3h). Yellow solid was obtained in 66% yield. Mp 89–90 °C; ^1H NMR (500 MHz, CDCl_3) δ 2.63–2.65 (m, 2H), 4.03–4.06 (m, 2H), 4.97 (d, $J = 2.0$ Hz, 2H), 6.52 (dd, $J = 2.0, 3.5$ Hz, 1H), 6.64 (d, $J = 3.5$ Hz, 1H), 7.38 (dd, $J = 2.0$ Hz, 1H), 7.58 (m, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 39.8, 65.5, 69.1, 113.1, 118.4, 121.8, 129.8, 146.2, 151.9, 195.9 ppm; IR (neat) ν 1675, 1595, 1263 cm^{-1} ; MS (70 eV): m/z 178 (M^+), 150, 122, 94. Anal. Calcd for $\text{C}_{10}\text{H}_{10}\text{O}_3$: C, 67.41, H, 5.66. Found: C 66.89; H, 5.55.

(E)-3-(Thiophen-2-ylmethylene)dihydro-2H-pyran-4(3H)-one (3i). Yellow solid was obtained in 65% yield. Mp 116–117 °C; ^1H NMR (500 MHz, CDCl_3) δ 2.64–2.67 (m, 2H), 4.05–4.07 (m, 2H), 4.86 (d, $J = 2.0$ Hz, 2H), 7.15 (dd, $J = 3.7, 5.0$ Hz, 1H), 7.27 (d, $J = 3.7$ Hz, 1H), 7.58 (d, $J = 5.0$ Hz, 1H), 7.82 (dd, $J = 2.0, 2.0$ Hz, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 39.7, 65.5, 68.9, 128.4, 128.7, 130.0, 131.9, 134.1, 138.2, 195.5 ppm; IR (neat) ν 2870, 1676, 1416 cm^{-1} ; MS (70 eV): m/z 194 (M^+), 166, 138, 110. Anal. Calcd for $\text{C}_{10}\text{H}_{10}\text{O}_2\text{S}$: C, 61.83; H, 5.19. Found: C, 61.78; H, 5.03.

(E)-3-(3-Methylbutylidene)dihydro-2H-pyran-4(3H)-one (3j). Yellow oil was obtained in 57% yield. ^1H NMR (500 MHz, CDCl_3) δ 0.89 (d, $J = 6.5$ Hz, 6H), 1.72–1.78 (m, 1H), 1.90 (dd, $J = 7.0, 7.5$ Hz, 2H), 2.53–2.56 (m, 2H), 3.95–3.98 (m, 2H), 4.51 (d, $J = 1.0$ Hz, 2H), 6.73–6.77 (m, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 22.9, 28.6, 36.9, 39.9, 65.7, 67.7, 134.1, 139.4, 196.0 ppm; IR (neat) ν 2958, 1696, 1466 cm^{-1} ; MS (70 eV): m/z 168 (M^+), 151, 112, 83.

(E)-1-Methyl-3-(2-methylbenzylidene)piperidin-4-one (4a). Yellow oil was obtained in 70% yield. ^1H NMR (500 MHz, CDCl_3) δ 2.28 (s, 3H), 2.37 (s, 3H), 2.64–2.68 (m, 2H), 2.78–2.81 (m, 2H), 3.47 (d, $J = 1.5$ Hz, 2H), 7.05 (d, $J = 7.5$ Hz, 1H), 7.15–7.26 (m, 3H), 7.65 (s, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 20.4, 39.6, 46.4, 53.5, 57.6, 125.9, 129.2, 129.3, 130.7, 133.6, 134.3, 135.4, 138.5, 198.1 ppm; IR (neat) ν 2942, 1687, 1608 cm^{-1} ; MS (70

eV): m/z 215 (M^+) 198, 186, 115. Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{NO}$: C, 78.10; H, 7.96. Found: C, 78.22; H, 8.03.

(E)-1-Methyl-3-(2-methoxybenzylidene)piperidin-4-one (4b). Yellow oil was obtained in 68% yield. ^1H NMR (500 MHz, CDCl_3): δ 2.43 (s, 3H), 2.68–2.69 (m, 2H), 2.84–2.86 (m, 2H), 3.59 (s, 2H), 3.82 (s, 3H), 6.89 (d, $J = 7.5$ Hz, 1H), 6.94 (dd, $J = 7.0, 7.5$ Hz, 1H), 7.13 (d, $J = 7.0$ Hz, 1H), 7.32 (dd, $J = 7.0, 7.5$ Hz, 1H), 7.79 (s, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 39.3, 46.2, 53.3, 55.9, 57.8, 107.5, 111.2, 120.4, 124.2, 130.6, 131.1, 132.8, 158.9, 197.7 ppm; IR (neat) ν 2942, 1685, 1610 cm^{-1} ; MS (70 eV): m/z 231 (M^+), 203, 188, 144, 124. Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$: C, 72.70; H, 7.41. Found: C, 72.88; H, 7.63.

(E)-1-Methyl-3-(3-methoxybenzylidene)piperidin-4-one (4c). Yellow oil was obtained in 71% yield. ^1H NMR (500 MHz, CDCl_3) δ 2.44 (s, 3H), 2.47–2.67 (m, 2H), 2.82–2.88 (m, 2H), 3.67 (s, 2H), 3.80 (s, 3H), 6.86 (s, 1H), 6.88–6.97 (m, 2H), 7.30 (dd, $J = 7.5, 8.0$ Hz, 1H), 7.53 (s, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 39.4, 46.5, 53.1, 55.7, 58.0, 115.0, 116.3, 123.1, 129.9, 133.5, 136.2, 136.5, 159.9, 197.9 ppm; IR (neat) ν 2940, 1686, 1599 cm^{-1} ; MS (70 eV): m/z 231 (M^+), 202, 174, 145. Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$: C, 72.70; H, 7.41. Found: C, 72.97; H, 7.61.

(E)-3-(Furan-2-ylmethylene)-1-methylpiperidin-4-one (4d). Yellow oil was obtained in 73% yield. ^1H NMR (500 MHz, CDCl_3) δ 2.41 (s, 3H), 2.54–2.57 (m, 2H), 2.70–2.72 (m, 2H), 3.70 (s, 2H), 6.43 (dd, $J = 1.5, 1.5$ Hz, 1H), 6.56 (d, $J = 3.0$ Hz, 1H), 7.26 (s, 1H), 7.50 (s, 1H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 39.2, 46.6, 52.7, 57.8, 112.9, 117.7, 122.2, 129.5, 145.7, 152.1, 197.3 ppm; IR (neat) ν 2944, 1681, 1598 cm^{-1} ; MS (70 eV): m/z 191 (M^+), 162, 134, 106. Anal. Calcd for $\text{C}_{11}\text{H}_{13}\text{NO}_2$: C, 69.09; H, 6.85. Found: C, 69.15; H, 6.72.

3. Results and discussion

We first optimized the conditions for the reaction of **1a** with benzaldehyde, as summarized in Table 1. The best results were obtained under solvent-free conditions and in the presence of $\text{MgBr}_2 \cdot \text{OEt}_2$ (7 mol%) and TMSNMe_2 (entry 1). Use of other amines, such as NHET_2 (entry 2) and NEt_3 (entry 3), gave lower yields of **3a** and, therefore, formation of more quantities of the respective bis compound was observed. Omission of either the Lewis acid (entry 4) or the amine (entry 5) from the mixture led to the nearly complete recovery of the starting materials, proving that both additives are

Table 1. Optimization of the reaction conditions for the synthesis of **3a**.

Entry	Lewis acid	Amine ^a	Time (h)	Yield (%) ^b
1	MgBr ₂ .OEt ₂	TMSNMe ₂	6	73
2	MgBr ₂ .OEt ₂	NHEt ₂	6	35
3	MgBr ₂ .OEt ₂	NEt ₃	6	27
4	–	TMSNMe ₂	12	< 5
5	MgBr ₂ .OEt ₂	–	12	< 5
6	Mg(ClO ₄) ₂	TMSNMe ₂	6	51
7	MgF ₂	TMSNMe ₂	6	39
8	LiClO ₄	TMSNMe ₂	6	44
9	LiBr	TMSNMe ₂	6	41


^a: 0.50 equivalents;^b: Isolated yields.

necessary for the progress of the reaction. Use of other Lewis acids could not also improve the yield (entries 6-9).

Under optimized conditions (MgBr₂.OEt₂/TMSNMe₂), other aldehydes possessing electron withdrawing and electron releasing substituents behaved similarly to produce **3b-g** within 3-7 h (Table 2, entries 1-6). Use of heteroaromatic aldehydes (entries 7-8) further showed the generality of the procedure. A moderate yield was obtained for the same reaction with 3-methylbutanal **2j** (entry 9), an example of aliphatic aldehydes, which are also prone to self-condensation and usually give very low yields of the condensation product under aldol conditions.

Generally, the above reactions proceeded at ambient temperature and reached completion in short time periods, as indicated in the tables. ¹H NMR analysis of the reaction mixtures suggested that a single monoarylidene isomer with E geometry for the olefinic bond was formed in each case as the major product of the reaction. Comparison of ¹H NMR spectra shows that in the products possessing *ortho* substituents on the aldehyde residue, the H-2 protons shift slightly to lower fields. In particular, in the case of **3b**, the *ortho* fluorine atom causes the C-2 carbon signal to appear as a doublet in the ¹³C NMR spectrum. Similarly, in the ¹H NMR spectrum, the C-2 protons, which are already coupled to the exocyclic olefinic proton, show an extra coupling with the fluorine atom, presumably due to a “through-space” hydrogen-fluorine interaction (Figure 1). These features provide solid evidence for the absolute stereochemistry of the exocyclic double bond.

To show the generality of the method, we then subjected ketone **1b** to the same conditions. Scheme 2 summarizes the results of the reactions of **1b** with



Product	Ar	Time (h)	Yield (%) ^a
4a	2-MeC ₆ H ₄	4	70
4b	2-MeOC ₆ H ₄	5	68
4c	3-MeOC ₆ H ₄	3	71
4d	4-ClC ₆ H ₄	3	70
4e	2-furyl	4	73

^a: Isolated yields.**Scheme 2.** The reaction pathway for the synthesis of **4**.

various aldehydes, leading to formation of products **4a-e**, within 3-5 h time periods.



















Based on these results, we propose a mechanism for the reaction (Figure 2). First, TMSNMe₂ facilitates conversion of the starting aldehyde into its respective in situ formed iminium salt. The trimethylsilanolate anion resulted from this conversion, then causes the enolization of the ketone **1** under the catalysis of MgBr₂.OEt₂. At last, the reaction of the enol with the iminium ion gives the final product. This kind of mechanism for similar aldol condensation reactions involving iminium ion intermediates has precedence [15,16].

4. Conclusion

A significant advantage of the present method is the observed chemoselectivity, which leads to minimum formation of the unwanted bis byproducts. This must be due to the relatively mild conditions chosen for this reaction. In other words, formation of a bis byproduct has to go through enolization of **3**. This enolization is less feasible in the presence of a mild amine such as TMSNMe₂, because the protons of the starting ketone **1** are expected to be more acidic than those of **3** (four protons and a simple C=O in **1** vs. two protons and a conjugated C=O in **3**). Further, use of equimolar amounts of the aldehyde itself exhibits extensive formation of the undesired bis compounds.

In summary, a convenient protocol is presented for the efficient synthesis of monoarylidenes of two synthetically important heterocyclic ketones. The procedure takes place in one pot and involves mild reaction conditions. Moreover, the workup is easy and the use of expensive and commercially unavailable reagents is avoided, while equimolar quantities of the starting materials and trace amounts of the catalyst are employed. A search in the literature shows that there are only scattered reports available on the synthesis of a few derivatives of the title compounds [20,30-32] and this makes the present procedure a valuable addition to the related archive.

Table 2. Room-temperature $\text{MgBr}_2 \cdot \text{OEt}_2$ catalyzed synthesis of **3**.

Entry	Aldehyde	Product	Time (h)	Yield% ^a
1	 2b	 3b	3	75
2	 2c	 3c	6	70
3	 2d	 3d	4	73
4	 2e	 3e	3	76
5	 2f	 3f	7	68
6	 2g	 3g	6	86
7	 2h	 3h	4	66
8	 2i	 3i	4	65
9	 2j	 3j	7	57

^a: Isolated yields.




Figure 1. Elucidative “through-space” C-F (top) and H-F (bottom) couplings in **3b**.




Figure 2. The suggested mechanism for the reaction.

Acknowledgement

Partial financial support by the Ministry of Science, Research, and Technology of Iran is gratefully appreciated.

References

- LoPachin, R.M., Barber, D.S. and Gavin, T. “Molecular mechanisms of the conjugated α,β -unsaturated carbonyl derivatives: Relevance to neurotoxicity and neurodegenerative diseases”, *Toxicol. Sci.*, **104**(2), pp. 235-249 (2008).
- Romanoa, J.J. and Casillas, E. “A short synthesis of morachalcone A”, *Tetrahedron Lett.*, **46**(13), pp. 2323-2326 (2005).
- Luo, Y., Song, R., Li, Y., Zhang, S., Liu, Z.-J., Fu, J. and Zhu, H.-L. “Design, synthesis, and biological evaluation of chalcone oxime derivatives as potential

- immunosuppressive agents" *Bioorg. Med. Chem. Lett.*, **22**(9), pp. 3039-3043 (2012).
4. Juvale, K., Pape, V.F.S. and Wiese, M. "Investigation of chalcones and benzochalcones as inhibitors of breast cancer resistance protein", *Bioorg. Med. Chem.*, **20**(1), pp. 346-355 (2012).
 5. Rodrigues, J.A.R., Moran, P.J.S., Conceição, G.J.A. and Fardelone, L.C. "Recent advances in the biocatalytic asymmetric reduction of acetophenones and α, β -unsaturated carbonyl compounds", *Food Technol. Biotechnol.*, **42**(4), pp. 295-303 (2004).
 6. Wang, Y., Ye, J. and Liang, X. "Convenient preparation of chiral α, β -epoxy ketones via Claisen-Schmidt condensation-epoxidation sequence", *Adv. Synth. Catal.*, **349**(7), pp. 1033-1036 (2007).
 7. Rajput, J.K. and Kaur, G. "Silicotungstic acid catalysed Claisen-Schmidt condensation reaction: An efficient protocol for synthesis of 1,3-diaryl-2-propenones", *Tetrahedron Lett.*, **53**(6), pp. 646-649 (2012).
 8. Nicolaou, K.C., Zhong, Y.-L. and Baran, P.S. "A new method for the one-step synthesis of α, β -unsaturated carbonyl systems from saturated alcohols and carbonyl compounds", *J. Am. Chem. Soc.*, **122**(31), pp. 7596-7597 (2000).
 9. Othman, M.R., Helwani, Z., Martunus and Fernando, W.J.N. "Synthetic hydrotalcites from different routes and their application as catalysts and gas adsorbents", *Appl. Organomet. Chem.*, **23**(9), pp. 335-346 (2009).
 10. Leonova, E., Makarov, M., Klemenkova, Z. and Odinet, I. "Lewis acids as mild and effective catalysts for the synthesis of 3,5-bis[(hetero)aryliden]piperidin-4-ones", *Helv. Chim. Acta*, **93**(10), pp. 1990-1999 (2010).
 11. Qiu, R., Qiu, Y., Yin, S., Xu, X., Luo, S., Au, C.-T., Wong, W.-Y. and Shimada, S. "Highly efficient and selective synthesis of (E)- α, β -unsaturated ketones by crossed condensation of ketones and aldehydes catalyzed by an air-stable cationic organobismuth perfluorooctanesulfonate", *Adv. Synth. Catal.*, **352**(1), pp. 153-162 (2010).
 12. Qiu, R., Qiu, Y., Yin, S., Song, X., Meng, Z., Xu, X., Zhang, X., Luo, S., Au, C.-T. and Wong, W.-Y. "Facile separation catalyst system: Direct diastereoselective synthesis of (E)- α, β -unsaturated ketones catalyzed by an air-stable Lewis acidic/basic bifunctional organobismuth complex in ionic liquids", *Green Chem.*, **12**(10), pp. 1767-1771 (2010).
 13. Wang, J.-F., Lei, M., Li, Q., Ge, Z.-m., Wang, X. and Li, R.-t. "A novel and efficient direct aldol condensation from ketones and aromatic aldehydes catalyzed by proline-TEA through a new pathway", *Tetrahedron*, **65**(25), pp. 4826-4833 (2009).
 14. Yang, S.-D., Wu, L.-Y., Yan, Z.-Y., Pan, Z.-L. and Liang, Y.-M. "A novel ionic liquid supported organocatalyst of pyrrolidine amide: Synthesis and catalyzed Claisen-Schmidt reaction", *J. Mol. Catal. A Chem.*, **268**(1-2), pp. 107-111 (2007).
 15. Kreher, U.P., Rosamilia, A.E., Raston, C.L., Scott, J.L. and Strauss, C.R. "Direct preparation of monoarylidene derivatives of aldehydes and enolizable ketones with DIMCARB", *Org. Lett.*, **5**(17), pp. 3107-3110 (2003).
 16. Wang, W., Mei, Y., Li, H. and Wang, J. "A novel pyrrolidine imide catalyzed direct formation of α, β -unsaturated ketones from unmodified ketones and aldehydes", *Org. Lett.*, **7**(4), pp. 601-604 (2005).
 17. Abaee, M.S., Mojtahedi, M.M., Hamidi, V., Mesbah, A.W. and Massa, W. "The first synthesis of bis(arylmethylidene)dioxan-5-ones: Potential scaffolds to access vicinal tricarbonyl derivatives", *Synthesis*, **40**(13), pp. 2122-2126 (2008).
 18. Abaee, M.S., Mojtahedi, M.M. and Zahedi, M.M. "An efficient and improved method for the synthesis of bis(arylmethylidene)thiopyranones", *Synlett*, **16**(15), pp. 2317-2320 (2005).
 19. Mojtahedi, M.M., Abaee, M.S., Samianifard, M., Shamloo, A., Padyab, M., Mesbah, A.W. and Harms, K. "Ultrasound mediation for efficient synthesis of monoarylidene derivatives of homo- and heterocyclic ketones", *Ultrason. Sonochem.*, **20**(3), pp. 924-930 (2013).
 20. Mojtahedi, M.M., Abaee, M.S., Khakbaz, M., Alishiri, T., Samianifard, M., Mesbah, A.W. and Harms, K. "An efficient procedure for the synthesis of α, β -unsaturated ketones and its application to heterocyclic systems", *Synthesis*, **43**(23), pp. 3821-3826 (2011).
 21. Mojtahedi, M.M., Abaee, M.S., Samianifard, M., Shamloo, A., Padyab, M., Mesbah, A.W. and Harms, K. "Ultrasound mediation for efficient synthesis of monoarylidene derivatives of homo- and heterocyclic ketones", *Ultrason. Sonochem.*, **20**(3), pp. 924-930 (2013).
 22. Leflemme, N., Dallemagne, P. and Rault, S. "An efficient route to 6-(het)aryl-2-methyl-2,3-dihydro-1H-pyridin-4-ones as potential nAChRs ligands", *Tetrahedron*, **60**(22), pp. 4861-4865 (2004).
 23. Watson, P.S., Jiang, B. and Scott, B. "A diastereoselective synthesis of 2,4-disubstituted piperidines: Scaffolds for drug discovery", *Org. Lett.*, **2**(23), pp. 3679-3681 (2000).
 24. Etayo, P., Badorrey, R., Díaz-de-Villegas, M.D. and Gálvez, J.A. "Stereoselective synthesis of chiral 2,3-disubstituted 2,3-dihydro-4(1H)-pyridones", *Eur. J. Org. Chem.*, **35**, pp. 6008-6014 (2008).
 25. Xavier, N.M. and Rauter, A.P. "Sugars containing α, β -unsaturated carbonyl systems: Synthesis and their usefulness as scaffolds in carbohydrate chemistry", *Carbohydr. Res.*, **343**(10-11), pp. 1523-1539 (2008).
 26. Clarke, P.A. and Santos, S. "Strategies for the formation of tetrahydropyran rings in the synthesis of natural products", *Eur. J. Org. Chem.*, **9**, pp. 2045-2053 (2006).

27. Pellissier, H. "Asymmetric hetero-Diels-Alder reactions of carbonyl compounds", *Tetrahedron*, **65**(15), pp. 2839-2877 (2009).
28. Lichtenthaler, F.W., Nakamura, K. and Klotz, J. "(-)-Daucic acid: Revision of configuration, synthesis, and biosynthetic implications", *Angew. Chem. Int. Ed.*, **42**(47), pp. 5838-5843 (2003).
29. Pohmakotr, N., Geiss, J.H. and Seebach, D. "Applications of magnesium thioacrolein dianions to the synthesis of sulfur-free compounds (vinyl- and divinylepoxides, acylcyclopentenones)" [Anwendungen der magnesiumderivate von thioacrolein-dianionen zur synthese schwefelfreier verbindungen (vinyl- und divinylepoxide, acylcyclopentene)], *Chem. Ber.*, **112**(4), pp. 1420-1439 (1979).
30. Liotta, F., Lu, H., Wachter, M.P. and Xia, M. "Tetrahydro-pyranonpyrazolecannabinoid modulator", US Patent 0069341 A1 (2010).
31. Kumar, R.R., Perumal, S., Senthilkumar, P., Yogeeswari, P. and Sriram, D. "A facile synthesis and antimycobacterial evaluation of novel spiro-pyrrolopyrrolizines and pyrrolidines", *Eur. J. Med. Chem.*, **44**(9), pp. 3821-3829 (2009).
32. Gayathri, D., Velmurugan, D., Kumar, R.R., Perumal, S. and Ravikumar, K. "3-[(E)-4-Methoxybenzylidene]-1-methylpiperidin-4-one", *Acta Cryst.*, E64, p. o552 (2008).

Biographies

Mohammad Majid Mojtahedi obtained BS and MS degrees from Isfahan University, Iran, and Shahid

Beheshti University, Iran, respectively, and a PhD degree from Sharif University of Technology, Tehran, Iran. He is currently Associate Professor in the Faculty of Organic Chemistry and Natural Products at the Chemistry and Chemical Engineering Research Center of Iran (CCERCI). His research interests include new methodologies in organic synthesis, green chemistry and medicinal chemistry.

Mohammad Saeed Abaee obtained his MS degree from the University of Tehran, Iran, and a PhD degree from the University of Saskatchewan, Canada, in 1999. He is currently Associate Professor in the Faculty of Organic Chemistry and Natural Products at the Chemistry and Chemical Engineering Research Center of Iran (CCERCI). His research interests include the study of Lewis acid catalyzed reactions, thiopyran heterocycles, and environmental chemistry.

Maedeh Samianifard obtained her BS degree from the Islamic Azad University, Iran, and is currently studying for her MS degree at the Chemistry and Chemical Engineering Research Center of Iran (CCERCI). Her research interests include new synthetic methodologies in organic chemistry.

Akram Shamloo obtained her BS degree from the Islamic Azad University, Iran, and is currently studying for her MS degree in the Chemistry and Chemical Engineering Research Center of Iran (CCERCI). Her research interests include new synthetic methodologies in organic chemistry.